

A DEGREE BOUND FOR GLOBALLY GENERATED VECTOR BUNDLES

JOSÉ CARLOS SIERRA

ABSTRACT. Let E be a globally generated vector bundle of rank $e \geq 2$ over a reduced irreducible projective variety X of dimension n defined over an algebraically closed field of characteristic zero. Let $L := \det(E)$. If $\deg(E) := \deg(L) = L^n > 0$ and E is not isomorphic to $\mathcal{O}_X^{e-1} \oplus L$, we obtain a sharp bound

$$\deg(E) \geq h^0(X, E) - e$$

on the degree of E , proving also that $\deg(L) = h^0(X, L) - n$ if equality holds. As an application, we obtain a Del Pezzo-Bertini type theorem on varieties of minimal degree for subvarieties of Grassmannians, as well as a bound on the sectional genus for subvarieties $X \subset \mathbb{G}(k, N)$ of degree at most $N + 1$.

1. INTRODUCTION

Let X be a reduced irreducible projective variety of dimension n defined over an algebraically closed field k of characteristic zero.

If $X \subset \mathbb{P}^N$ is a non-degenerate embedding (i.e. not contained in any hyperplane) then

$$\deg(X) \geq N + 1 - n,$$

and $X \subset \mathbb{P}^N$ is said to be a variety of minimal degree if equality holds. These varieties were classified by Del Pezzo [2] for $n = 2$, and Bertini [1] for $n \geq 3$ (see also [3] for a modern account). A variety of minimal degree turns out to be either a linear space, or a quadric hypersurface, or a rational normal scroll, or (a cone over) the Veronese surface in $\mathbb{P}^5$.

More generally, if L is a globally generated line bundle on X and $\deg(L) := L^n > 0$, then

$$\deg(L) \geq h^0(X, L) - n,$$

and, if equality holds, then $\varphi_{|L|} : X \rightarrow X' \subset \mathbb{P}^{h^0(X, L)-1}$ is a birational morphism onto a variety of minimal degree, so in particular X is rational.

In this note, a similar bound is obtained for a globally generated vector bundle E of rank $e \geq 2$ over X such that $\deg(E) := \deg(L) = L^n > 0$, where $L := \det(E)$. If $E \cong \mathcal{O}_X^{e-1} \oplus L$ then

$$\deg(E) = \deg(L) \geq h^0(X, L) - n = h^0(X, E) - e + 1 - n.$$

Therefore, we assume E different from $\mathcal{O}_X^{e-1} \oplus L$ in the sequel. The main result of the paper is the following:

Theorem 1. *Let E be a globally generated vector bundle of rank $e \geq 2$ on X and let $L := \det(E)$. If $\deg(E) > 0$ and E is not isomorphic to $\mathcal{O}_X^{e-1} \oplus L$, then*

$$\deg(E) \geq h^0(X, E) - e.$$

Research partially supported by MCYT project MTM2006-04785 and by the program “Profesores de la UCM en el extranjero. Convocatoria 2006”.

2000 *Mathematics Subject Classification*: Primary 14F05, 14N25. Secondary 14M15.

Moreover, if equality holds then $\deg(L) = h^0(X, L) - n$ and, in particular, X is rational.

We remark that this bound does not depend on the dimension of X , as one might expect a priori.

For general Cartier divisors $Y_1, \dots, Y_{n-1} \in |L|$, let $C := \cap_{i=1}^{n-1} Y_i$ denote the curve section (consider $C = X$ for $n = 1$) and let $p_a(C)$ denote its arithmetic genus. As a byproduct of the proof of Theorem 1, we also obtain the following result:

Corollary 1. *Let E be a globally generated vector bundle of rank $e \geq 2$ on X and let E^* denote its dual bundle. Assume $h^0(X, E) > e + 1$ for $n \geq 2$. If $\deg(E) > 0$ and E is not isomorphic to $\mathcal{O}_X^{e-1} \oplus L$, then:*

- (i) *if $\deg(E) < h^0(X, E) - h^0(X, E^*)$ then $C \cong \mathbb{P}^1$. Moreover, if X is normal then $\deg(L) = h^0(X, L) - n$ and X is rational;*
- (ii) *if $\deg(E) = h^0(X, E) - h^0(X, E^*)$ then either $p_a(C) \leq 1$, or C is hyperelliptic and the restriction of E to C satisfies $E_C \cong \mathcal{O}_C^r \oplus g_2^1 \oplus \dots \oplus g_2^1$, where $r := h^0(C, E_C^*)$ and g_2^1 denotes the pencil of degree 2 on C .*

Remark 1. For $n = 1$, both Theorem 1 and Corollary 1 hold under weaker hypotheses (see Lemma 2 and Proposition 2).

As an application of Theorem 1, we extend Del Pezzo-Bertini's theorem to non-degenerate subvarieties $X \subset \mathbb{G}(k, N)$ of Grassmannians of k -planes in $\mathbb{P}^N$. In this context, *non-degenerate* means that X is not contained in any $\mathbb{G}(k, N - 1)$:

Corollary 2. *Let $X \subset \mathbb{G}(k, N)$ be a non-degenerate embedding. If $X \subset \mathbb{G}(k, N)$ is not contained in any Schubert variety $\Omega(k - 1, N)$ of k -planes containing a linear subspace $\mathbb{P}^{k-1} \subset \mathbb{P}^N$ then*

$$\deg(X) \geq N - k,$$

where $\deg(X)$ is the degree of $X \subset \mathbb{P}^M$ given by the Plücker embedding of $\mathbb{G}(k, N)$. Moreover, if equality holds then $X \subset \mathbb{P}^M$ is a variety of minimal degree in its linear span.

Furthermore, we can also rephrase Corollary 1 for subvarieties of Grassmannians. Let $X \subset \mathbb{G}(k, N)$, and let $\mathcal{O}_X(1)$ denote the line bundle on X giving the Plücker embedding $X \subset \mathbb{P}^M$. For general Cartier divisors $H_1, \dots, H_{n-1} \in \mathcal{O}_X(1)$, let $C := \cap_{i=1}^{n-1} H_i$ denote the curve section (consider $C = X$ for $n = 1$) and let $p_a(C)$ denote its arithmetic genus:

Corollary 3. *Let $X \subset \mathbb{G}(k, N)$ be a non-degenerate embedding. Assume $N - k \geq 2$ for $n \geq 2$. Let $s \leq k - 1$ be a non-negative integer. If $X \subset \mathbb{G}(k, N)$ is not contained in any Schubert variety $\Omega(s, N)$ of k -planes containing a linear subspace $\mathbb{P}^s \subset \mathbb{P}^N$, then:*

- (i) *if $\deg(X) \leq N - s$ then $C \cong \mathbb{P}^1$. Moreover, if X is normal then $X \subset \mathbb{P}^M$ is (a linear projection of) a variety of minimal degree in its linear span;*
- (ii) *if $\deg(X) = N + 1 - s$ then $p_a(C) \leq 1$.*

It is proved in [7] that non-degenerate manifolds $X \subset \mathbb{P}^N$ of degree $\deg(X) \leq N$ are simply connected. This topological result follows as a corollary of a complete classification of such manifolds. According to Corollary 3, the Plücker embedding of a non-degenerate normal subvariety $X \subset \mathbb{G}(k, N)$ of degree $\deg(X) \leq N$ not contained in any Schubert variety $\Omega(0, N)$ is either a linear space, or a quadric hypersurface, or a rational scroll, or (a cone over) the Veronese surface. Hence, we also deduce *a posteriori* a similar result for subvarieties of Grassmannians:

Corollary 4. *Assume $N - k \geq 2$ for $n \geq 2$. Then, a non-degenerate normal subvariety $X \subset \mathbb{G}(k, N)$ of degree $\deg(X) \leq N$ not contained in any Schubert variety $\Omega(0, N)$ is simply connected.*

The bound $\deg(X) \leq N$ is optimal in both [7] and Corollary 4. Indeed, for any $r \geq 1$ there exist r -dimensional elliptic scrolls in $\mathbb{P}^{2r}$ of degree $2r + 1$. Considering the corresponding elliptic curve in $\mathbb{G}(r - 1, 2r)$ of degree $2r + 1$, we obtain that also our bound is sharp. On the other hand, if we consider a cone in $\mathbb{P}^{2r+1}$ of vertex a point over an r -dimensional elliptic scroll in $\mathbb{P}^{2r}$ of degree $2r + 1$, the corresponding elliptic curve of degree $2r + 1$ in $\mathbb{G}(r, 2r + 1)$ shows that the assumption X not contained in any Schubert variety $\Omega(0, N)$ cannot be dropped.

The paper is organized as follows. Section 2 is devoted to the proof of Theorem 1 and Corollary 1. To this purpose, the crucial point here is to prove that $h^0(X, E) \leq h^0(C, E_C)$ whenever E is not isomorphic to $\mathcal{O}_X^{e-1} \oplus L$ and $\deg(E) > 0$ (see Corollary 5). Then we obtain a degree bound for globally generated vector bundles on curves (see Lemma 2 and Proposition 2). Using the correspondence between globally generated vector bundles and maps to Grassmannians, in Section 3 we translate Theorem 1 and Corollary 1 into Corollary 2 and Corollary 3, respectively. Some examples of subvarieties on the boundary are also included.

2. PROOF OF THEOREM 1

Let E be a globally generated vector bundle of rank $e \geq 2$ on X and let $L := \det(E)$.

Proposition 1. *If $h^0(X, E \otimes L^{-1}) \neq 0$ then $E \cong \mathcal{O}_X^{e-1} \oplus L$.*

Proof. Let $s \in H^0(X, E \otimes L^{-1})$ be a non-zero section. Consider the corresponding exact sequence of sheaves on X

$$0 \longrightarrow L \longrightarrow E \longrightarrow F \longrightarrow 0.$$

Let $Z = \{x \in X \mid s(x) = 0\}$ denote the zero-locus of s . We claim that $Z = \emptyset$. To get a contradiction, let $x \in Z$ and let $C \subset X$ be a curve not contained in Z such that $x \in C$. Restricting the above exact sequence to the normalization $\tilde{C}$ of C , we obtain an exact sequence of sheaves

$$0 \longrightarrow L_{\tilde{C}} \longrightarrow E_{\tilde{C}} \longrightarrow Q \oplus t \longrightarrow 0,$$

where Q is a vector bundle of rank $e - 1$ on $\tilde{C}$ and t is a torsion sheaf on $\tilde{C}$. Then

$$\deg(Q) + \text{length}(t) = \deg(E_{\tilde{C}}) - \deg(L_{\tilde{C}}) = 0.$$

This yields a contradiction since Q is globally generated, so $\deg(Q) \geq 0$, and $x \in \text{Supp}(t)$, so $\text{length}(t) \geq 1$.

We deduce from $Z = \emptyset$ that F is a globally generated vector bundle of rank $e - 1$ over X . Thus there exists a morphism of vector bundles $u : \mathcal{O}_X^{e-1} \rightarrow F$ of rank $e - 1$ at some point $x \in X$, and taking determinants we get a morphism $\det(u) : \mathcal{O}_X \rightarrow \det(F)$. Then u has rank $e - 1$ at every $x \in X$, since $\det(F) \cong c_1(F) \cong c_1(E) - c_1(L) \cong \mathcal{O}_X$. Therefore $F \cong \mathcal{O}_X^{e-1}$, whence $E \cong \mathcal{O}_X^{e-1} \oplus L$ since E is globally generated (cf. Remark 2). $\square$

Remark 2. Let E^* denote the dual bundle of E . If E is a globally generated vector bundle on X we recall that $h^0(X, E^*) = r$ if and only if $E \cong \mathcal{O}_X^r \oplus F$, where F is a globally generated vector bundle with $h^0(X, F^*) = 0$. Moreover, if X is a curve, then a globally generated vector bundle E on X is ample if and only if $h^0(X, E^*) = 0$.

For any Cartier divisor $Y \in |L|$, let E_Y denote the restriction of E to Y . It follows from Remark 2 that $h^0(Y, E_Y^*) \geq h^0(X, E^*)$. Moreover:

Lemma 1. *If $h^0(Y, E_Y^*) > h^0(X, E^*)$ for some $Y \in |L|$, then $h^0(X, E) = e + 1$.*

Proof. Let $s := h^0(X, E)$. Consider the exact sequence of vector bundles

$$0 \longrightarrow G \longrightarrow \mathcal{O}_X^s \longrightarrow E \longrightarrow 0,$$

where G arises as the kernel of the evaluation morphism. Dualizing the exact sequence above and restricting to Y , we obtain the following commutative diagram:

$$\begin{array}{ccccccc} & & 0 & & 0 & & 0 \\ & & \downarrow & & \downarrow & & \downarrow \\ 0 & \longrightarrow & E^* \otimes L^{-1} & \longrightarrow & \mathcal{O}_X^s \otimes L^{-1} & \longrightarrow & G^* \otimes L^{-1} \longrightarrow 0 \\ & & \downarrow & & \downarrow & & \downarrow \\ 0 & \longrightarrow & E^* & \longrightarrow & \mathcal{O}_X^s & \longrightarrow & G^* \longrightarrow 0 \\ & & \downarrow & & \downarrow & & \downarrow \\ 0 & \longrightarrow & E_Y^* & \longrightarrow & \mathcal{O}_Y^s & \longrightarrow & G_Y^* \longrightarrow 0 \\ & & \downarrow & & \downarrow & & \downarrow \\ & & 0 & & 0 & & 0 \end{array}$$

If $h^0(Y, E_Y^*) > h^0(X, E^*)$, then a simple diagram chase shows that $h^0(X, G^* \otimes L^{-1}) \neq 0$. It follows from Proposition 1 that $G^* \cong \mathcal{O}_X^{s-e-1} \oplus L$, where $L = \det(E) \cong \det(G^*)$. In particular $G \cong \mathcal{O}_X^{s-e-1} \oplus L^{-1}$, whence $s = e + 1$ since $h^0(X, G) = 0$. $\square$

Assume furthermore $n \geq 2$ and $\deg(E) > 0$. For general Cartier divisors $Y_1, \dots, Y_{n-1} \in |L|$, let $C := \cap_{i=1}^{n-1} Y_i$ denote the curve section. Proposition 1 and Lemma 1 yield the following:

Corollary 5. *Assume $n \geq 2$ and $\deg(E) > 0$. If E is not isomorphic to $\mathcal{O}_X^{e-1} \oplus L$, then:*

- (i) $h^0(X, E) \leq h^0(C, E_C)$;
- (ii) if $h^0(X, E) > e + 1$, then $h^0(X, E^*) = h^0(C, E_C^*)$.

Proof. We first remark that $\cap_{i=1}^j Y_i$ is reduced and irreducible for every $j \in \{1, \dots, n-1\}$. This follows from Bertini's theorems applied to the morphism $\varphi_{|L|} : X \rightarrow \mathbb{P}^{h^0(X, L)-1}$ (see, for instance, [8, Corollary 6.11]). Here the hypothesis $\text{char}(k) = 0$ is used. If $h^0(X, E) = e + 1$ then $h^0(C, E_C) \geq e + 1$, since E_C is globally generated and non-trivial. Assume now $h^0(X, E) > e + 1$. Consider the exact sequence

$$0 \longrightarrow E \otimes L^{-1} \longrightarrow E \longrightarrow E_{Y_1} \longrightarrow 0.$$

It follows from Proposition 1 that $h^0(X, E \otimes L^{-1}) = 0$. Hence $h^0(X, E) \leq h^0(Y_1, E_{Y_1})$. On the other side, Lemma 1 yields $h^0(X, E^*) = h^0(Y_1, E_{Y_1}^*)$. In particular, E_{Y_1} cannot be isomorphic to $\mathcal{O}_{Y_1}^{e-1} \oplus L_{Y_1}$. Combining Proposition 1 and Lemma 1 we recursively obtain $h^0(X, E) \leq h^0(Y_1, E_{Y_1}) \leq h^0(Y_1 \cap Y_2, E_{Y_1 \cap Y_2}) \leq \dots \leq h^0(C, E_C)$, and also $h^0(X, E^*) = h^0(Y_1, E_{Y_1}^*) = h^0(Y_1 \cap Y_2, E_{Y_1 \cap Y_2}^*) = \dots = h^0(C, E_C^*)$. $\square$

We now consider globally generated vector bundles on curves. The following lemma generalizes [7, Proposition 2]:

Lemma 2. *Let C be a reduced irreducible curve of arithmetic genus $p_a(C) \geq 1$. Let E be a globally generated vector bundle of rank $e \geq 1$ over C , and let E^* denote its dual bundle. Then*

$$\deg(E) \geq h^0(C, E) - h^0(C, E^*).$$

Proof. By Remark 2, we may assume that E is ample, and, in that case, it is enough to prove that $\deg(E) \geq h^0(C, E)$. We proceed by induction, as in [7]. Assume $e = 1$. If $h^1(C, E) = 0$ the result follows from Riemann-Roch's theorem for singular curves (see, for instance, [5, Example 18.3.4 (a)]). If $h^1(C, E) \neq 0$ then $\deg(E) \geq 2(h^0(C, E) - 1)$ by a generalization of Clifford's theorem (see [4, Theorem A]), so $\deg(E) \geq h^0(C, E)$ since $h^0(C, E) \geq 2$ unless $E \cong \mathcal{O}_C$. Suppose now $e \geq 2$. If $h^0(C, E) \leq e$ then $E \cong \mathcal{O}_C^e$ since E is globally generated, so we can assume $h^0(C, E) \geq e + 1$. So, if $p \in C$ is a general point, a general section $s \in H^0(C, E(-p))$ does not vanish at any singular point of C . Thus, it induces an exact sequence

$$0 \longrightarrow M \longrightarrow E \longrightarrow E' \longrightarrow 0,$$

where M is a line bundle of positive degree and E' is an ample vector bundle of rank $e - 1$ over C . Thus $\deg(E') \geq h^0(C, E')$ by the induction hypothesis, and using again Riemann-Roch's theorem as well as [4, Theorem A] it follows that $\deg(M) \geq h^0(C, M)$. These two facts together yield

$$\deg(E) = \deg(E') + \deg(M) \geq h^0(C, E') + h^0(C, M),$$

and then $\deg(E) \geq h^0(C, E)$, since $h^0(C, E') \geq h^0(C, E) - h^0(C, M)$. $\square$

Furthermore, we state separately the following result that may also be useful for some other purpose:

Proposition 2. *If equality holds in Lemma 2, then either $E \cong \mathcal{O}_C^e$, or $p_a(C) = 1$, or C is a hyperelliptic curve and $E \cong \mathcal{O}_C^r \oplus g_2^1 \oplus \cdots \oplus g_2^1$, where $r := h^0(C, E^*)$ and g_2^1 denotes the pencil of degree 2 on C .*

Proof. According to Remark 2, we can assume $h^0(C, E^*) = 0$. We proceed by induction on e . Assume $e = 1$. If $h^1(C, E) = 0$ then $p_a(C) = 1$ by Riemann-Roch's theorem. Suppose now $h^1(C, E) \neq 0$. Then $h^0(C, E) = \deg(E) \geq 2(h^0(C, E) - 1)$ by [4, Theorem A], whence $h^0(C, E) = 2$ and equality holds. Therefore C is hyperelliptic and $E \cong g_2^1$. If $e \geq 2$ and $\deg(E) = h^0(C, E)$, repeating the argument of Lemma 2, it follows that necessarily $\deg(M) = h^0(C, M)$ and $\deg(E') = h^0(C, E')$. The induction hypothesis yields that either $p_a(C) = 1$, or C is a hyperelliptic curve and $E' \cong g_2^1 \oplus \cdots \oplus g_2^1$. Moreover, in the latter case [4, Theorem A] implies $M \cong g_2^1$. Therefore E is an extension of $g_2^1 \oplus \cdots \oplus g_2^1$ by g_2^1 , and $h^0(C, E) = 2e$. Let us show that $E \cong g_2^1 \oplus \cdots \oplus g_2^1$. To the contrary, assume that E corresponds to a non-trivial extension $\xi \in \text{Ext}^1(g_2^1 \oplus \cdots \oplus g_2^1, g_2^1) \cong \text{Ext}^1(\mathcal{O}_C^{e-1}, \mathcal{O}_C) \cong H^1(C, \mathcal{O}_C)^{e-1}$. Then, there exists also a non-trivial extension $\xi' \in \text{Ext}^1(g_2^1, g_2^1) \cong H^1(C, \mathcal{O}_C)$, yielding a

commutative diagram:

$$\begin{array}{ccccccc}
 & & 0 & & 0 & & \\
 & & \downarrow & & \downarrow & & \\
 & & \oplus^{e-2} g_2^1 & \xlongequal{\quad} & \oplus^{e-2} g_2^1 & & \\
 & & \downarrow & & \downarrow & & \\
 0 & \longrightarrow & g_2^1 & \longrightarrow & E & \longrightarrow & \oplus^{e-1} g_2^1 \longrightarrow 0 \\
 & & \parallel & & \downarrow & & \downarrow \\
 0 & \longrightarrow & g_2^1 & \longrightarrow & F & \longrightarrow & g_2^1 \longrightarrow 0 \\
 & & \downarrow & & \downarrow & & \downarrow \\
 & & 0 & & 0 & & 0
 \end{array}$$

To get a contradiction, we prove that $h^0(C, E) < 2e$. To this aim it is enough to check that $h^0(C, F) < 4$. Let $D \in g_2^1$ and consider the corresponding commutative diagram:

$$\begin{array}{ccccccc}
 & & 0 & & 0 & & 0 \\
 & & \downarrow & & \downarrow & & \downarrow \\
 0 & \longrightarrow & \mathcal{O}_C & \longrightarrow & F(-D) & \longrightarrow & \mathcal{O}_C \longrightarrow 0 \\
 & & \downarrow & & \downarrow & & \downarrow \\
 0 & \longrightarrow & g_2^1 & \longrightarrow & F & \longrightarrow & g_2^1 \longrightarrow 0 \\
 & & \downarrow & & \downarrow & & \downarrow \\
 0 & \longrightarrow & g_2^1 \otimes \mathcal{O}_D & \longrightarrow & F \otimes \mathcal{O}_D & \longrightarrow & g_2^1 \otimes \mathcal{O}_D \longrightarrow 0 \\
 & & \downarrow & & \downarrow & & \downarrow \\
 & & 0 & & 0 & & 0
 \end{array}$$

Looking at the long exact sequences of the first two rows we obtain the following square $\alpha, \beta, \gamma, \delta$:

$$\begin{array}{ccccccc}
 0 & \longrightarrow & H^0(C, \mathcal{O}_C) & \longrightarrow & H^0(C, F(-D)) & \longrightarrow & H^0(C, \mathcal{O}_C) \xrightarrow{\alpha} H^1(C, \mathcal{O}_C) \\
 & & \downarrow & & \downarrow & & \downarrow \gamma \\
 0 & \longrightarrow & H^0(C, g_2^1) & \longrightarrow & H^0(C, F) & \longrightarrow & H^0(C, g_2^1) \xrightarrow{\beta} H^1(C, g_2^1)
 \end{array}$$

Then $\alpha(1) = \xi' \neq 0$ and $\delta(\xi') \neq 0$ for a suitable choice of $D \in g_2^1$, whence β is not identically zero. Therefore $h^0(C, F) < 4$ and we get a contradiction. $\square$

We now obtain our main result:

Proof of Theorem 1. Using the notation in Corollary 5, assume first that $p_a(C) \geq 1$. From Lemma 2, Remark 2 and Corollary 5 we get:

$$\deg(E) = \deg(E_C) \geq h^0(C, E_C) - h^0(C, E_C^*) \geq h^0(C, E_C) - e \geq h^0(X, E) - e.$$

Moreover, $\deg(E) > h^0(X, E) - e$, since $h^0(C, E_C^*) = e$ implies that E_C is trivial, so $\deg(E) = 0$. Assume now $p_a(C) = 0$, so $C \cong \mathbb{P}^1$. Corollary 5 and Riemann-Roch's

theorem give:

$$h^0(X, E) \leq h^0(C, E_C) = \deg(E_C) + e = \deg(E) + e,$$

proving the first part.

We have seen that if $\deg(E) = h^0(X, E) - e$, then necessarily $C \cong \mathbb{P}^1$ and $h^0(X, E) = h^0(C, E_C)$. Consider the morphism $\varphi : X \rightarrow \mathbb{G}(k, N)$ given by E , where $k = e - 1$ and $N = h^0(X, E) - 1$ (cf. Section 3). Let $\mathbb{G}(k, N) \subset \mathbb{P}^M$ denote the Plücker embedding. As E_C is spanned, there is an exact sequence:

$$0 \longrightarrow \mathcal{O}_C^{e-1} \longrightarrow E_C \longrightarrow L_C \longrightarrow 0.$$

Taking cohomology we deduce that the canonical map $\bigwedge^e H^0(C, E_C) \rightarrow H^0(C, L_C)$ is surjective. Moreover, as we have seen, the restriction map $H^0(X, E) \rightarrow H^0(C, E_C)$ is an isomorphism. It follows that the composition of the restriction $\varphi|_C$ and the Plücker embedding is given by the complete linear system $|L_C|$. So, $\varphi(C) \subset \mathbb{P}^M$ is a rational normal curve in its linear span. We have $\deg(L_C) = \deg(E_C) = h^0(C, E_C) - e = N - k$, whence the linear span of $\varphi(C) \subset \mathbb{P}^M$ is a $\mathbb{P}^{N-k}$ and therefore the linear span of $\varphi(X) \subset \mathbb{P}^M$ is a $\mathbb{P}^{N-k+n-1}$. Thus $h^0(X, L) \geq N - k + n = \deg(E) + n = \deg(L) + n$. So we get $0 \geq \deg(L) + n - h^0(X, L) \geq 0$, and therefore $\deg(L) = h^0(X, L) - n$. In this case, the corresponding map $\varphi|_{L|} : X \rightarrow X' \subset \mathbb{P}^{h^0(X, L)-1}$ is a birational morphism onto a variety of minimal degree, so X is rational. $\square$

We proceed with the proof of Corollary 1:

Proof of Corollary 1. Corollary 5 yields $h^0(X, E^*) = h^0(C, E_C^*)$, since we assume $h^0(X, E) > e + 1$ whenever $n \geq 2$. Let us prove (i). It follows from Corollary 5 that

$$\deg(E_C) = \deg(E) < h^0(X, E) - h^0(X, E^*) \leq h^0(C, E_C) - h^0(C, E_C^*).$$

Then $p_a(C) = 0$ by Lemma 2, and so $C \cong \mathbb{P}^1$. If, moreover, X is normal then it easily follows from [9, Theorem 2] that $\deg(L) = h^0(X, L) - n$. Hence X is rational as before. We now prove (ii). If $\deg(E) = h^0(X, E) - h^0(X, E^*)$ then $\deg(E_C) \leq h^0(C, E_C) - h^0(C, E_C^*)$, and the thesis follows from Lemma 2 and Proposition 2. $\square$

We remark that the assumption $h^0(X, E) > e + 1$ in Corollary 1 cannot be dropped (cf. Remark 5).

3. A DEGREE BOUND FOR SUBVARIETIES OF GRASSMANNIANS

We first recall the well-known correspondence between globally generated vector bundles and maps to Grassmannians (see, for instance, [6] p. 207).

If E is a globally generated vector bundle of rank $e = k + 1$ over an algebraic variety X and $V \subset H^0(X, E)$ is a linear subspace of dimension $N + 1$, any epimorphism $V \otimes \mathcal{O}_X \rightarrow E \rightarrow 0$ determines a morphism $\varphi : X \rightarrow \mathbb{G}(k, N)$ such that $\varphi(X)$ is not contained in any $\mathbb{G}(k, N - 1)$. Conversely, every morphism $\varphi : X \rightarrow \mathbb{G}(k, N)$ such that $\varphi(X)$ is not contained in any $\mathbb{G}(k, N - 1)$ corresponds to a globally generated vector bundle E of rank $k + 1$ on X and an epimorphism $V \otimes \mathcal{O}_X \rightarrow E \rightarrow 0$. In both cases $E \cong \varphi^*(Q_W)$, where Q denotes the universal quotient bundle of rank $k + 1$ on $\mathbb{G}(k, N)$ and Q_W denotes its restriction to $W := \varphi(X) \subset \mathbb{G}(k, N)$. We simply denote Q_X if φ is the inclusion. In particular, for $e = 1$ we recover the correspondence between globally generated line bundles and maps to projective spaces. Moreover, if $\mathbb{G}(k, N) \subset \mathbb{P}^M$ denotes the Plücker embedding of the Grassmannian then its restriction $W \subset \mathbb{P}^M$ is given by a base point free linear system contained in $|\mathcal{O}_W(1)|$, where $\mathcal{O}_W(1) := \det(Q_W)$.

Remark 3. Let $\Omega(s, N) \subset \mathbb{G}(k, N)$ denote the Schubert variety of k -planes containing a linear subspace $\mathbb{P}^s \subset \mathbb{P}^N$. Then, $X \subset \mathbb{G}(k, N)$ is contained in a $\Omega(s, N)$ if and only if Q_X has a trivial quotient $\mathcal{O}_X^{s+1}$, and this happens if and only if $E \cong \mathcal{O}_X^{s+1} \oplus F$, where F is a globally generated vector bundle on X of rank $k - s$ (cf. Remark 2).

Theorem 1 allows us to extend Del Pezzo-Bertini's theorem on varieties of minimal degree to subvarieties of Grassmannians:

Proof of Corollary 2. We deduce from Remark 3 and Theorem 1 that

$$\deg(X) = \deg(Q_X) \geq h^0(X, Q_X) - e,$$

so $\deg(X) \geq N - k$ since $h^0(X, Q_X) \geq N + 1$ and $e = k + 1$. Furthermore, if $\deg(X) = N - k$ the linear span of $X \subset \mathbb{P}^M$ is a $\mathbb{P}^{N-k+n-1}$, as we concluded at the end of the proof of Theorem 1. Therefore $X \subset \mathbb{P}^M$ is a variety of minimal degree in its linear span. $\square$

Remark 4. If $X \subset \mathbb{G}(k, N)$ is a non-degenerate embedding contained in a Schubert variety $\Omega(k - 1, N)$ or, equivalently, $Q_X \cong \mathcal{O}_X^k \oplus \mathcal{O}_X(1)$, then $X \subset \Omega(k - 1, N) = \mathbb{P}^{N-k} \subset \mathbb{P}^M$ is a non-degenerate embedding of X in $\mathbb{P}^{N-k}$, and hence the bound $\deg(X) \geq N - k + 1 - n$ cannot be improved.

Example 1. Let us show some examples of subvarieties $X \subset \mathbb{G}(k, N)$ of degree $\deg(X) = N - k$ and not contained in any $\Omega(k - 1, N)$:

- (i) Let $X = \mathbb{P}^n$. Consider the embedding $X = \mathbb{G}(n - 1, n)$ given by $E = T_{\mathbb{P}^n}(-1)$.
- (ii) Consider $X = \mathbb{G}(1, 3)$, which is a hyperquadric in $\mathbb{P}^5$ by the Plücker embedding.
- (iii) Let $X = \mathbb{P}^2$. Consider the embedding $X \subset \mathbb{G}(1, 5)$ given by $E = \mathcal{O}_{\mathbb{P}^2}(1) \oplus \mathcal{O}_{\mathbb{P}^2}(1)$. The Plücker embedding of X is a Veronese surface in $\mathbb{P}^5$.
- (iv) Consider a Schubert variety $X = \Omega(1, N) \subset \mathbb{G}(1, N)$. The Plücker embedding of X is a cone of vertex a point over the Segre embedding $\mathbb{P}^1 \times \mathbb{P}^{N-2}$, so $\deg(X) = N - 1$.

We proceed with the proof of Corollary 3:

Proof of Corollary 3. If $X \subset \mathbb{G}(k, N)$ is not contained in any $\Omega(s, N)$ then it follows from Remark 3 that $h^0(X, Q_X^*) \leq s$. Therefore

$$\deg(Q_X) = \deg(X) \leq N + 1 - s \leq h^0(X, Q_X) - h^0(X, Q_X^*).$$

We deduce from Corollary 1 that $p_a(C) \leq 1$ unless C is hyperelliptic and $Q_C \cong \mathcal{O}_C^r \oplus g_2^1 \oplus \cdots \oplus g_2^1$. But the latter cannot occur since the morphism $\varphi : C \rightarrow \mathbb{G}(k, N)$ corresponding to Q_C is not an embedding. Indeed, $\varphi : C \rightarrow \varphi(C) \subset \mathbb{P}^M$ is a double covering of a rational normal curve of degree $k + 1 - r$. $\square$

Remark 5. If $X \subset \mathbb{G}(k, k + 1) = \mathbb{P}^{k+1}$ then Corollary 3 does not hold, since the assumption $h^0(X, Q_X) > e + 1$ in Corollary 1 is not satisfied.

Example 2. Some examples of Del Pezzo manifolds in $\mathbb{G}(k, N)$ of degree $N + 1$ and not contained in any $\Omega(0, N)$, thus on the boundary of Corollary 3, are given by the intersection of $\mathbb{G}(1, 3) \subset \mathbb{P}^5$ with a general hyperquadric of $\mathbb{P}^5$, or by $X = \mathbb{G}(1, 4)$.

Acknowledgements. This research was begun while I was visiting the Mathematics Department at Università degli Studi Roma Tre in the fall of 2006. I would like to thank Angelo F. Lopez for his warm hospitality during the three months I spent there, as well as Mike Roth for pointing out the simple proof of Proposition 1. Finally, I would especially like to thank the referee for many helpful comments and suggestions.

REFERENCES

- [1] Bertini, E.: *Introduzione alla Geometria Proiettiva degli Iperspazi*. Pisa, Spoerri (1907)
- [2] Del Pezzo, P.: *Sulle superficie di ordine n immerse nello spazio di $n+1$ dimensioni*. *Rend. R. Acc. delle Scienze Fisiche e Mat. di Napoli* **24** 212–216 (1885)
- [3] Eisenbud, D., Harris, J.: *On varieties of minimal degree (a centennial account)*. *Algebraic geometry, Bowdoin, 1985* (Brunswick, Maine, 1985), 3–13. *Proc. Sympos. Pure Math.*, 46, Part 1, Amer. Math. Soc., Providence, RI (1987)
- [4] Eisenbud, D., Koh, J., Stillman, M.: *Determinantal equations for curves of high degree*. Appendix (with J. Harris). *Amer. J. Math.* **110**(3), 513–539 (1988)
- [5] Fulton, W.: *Intersection theory*. *Ergebnisse der Mathematik und ihrer Grenzgebiete* (3), 2. Springer-Verlag, Berlin (1984)
- [6] Griffiths, Ph., Harris, J.: *Principles of algebraic geometry*. *Pure and Applied Mathematics*. Wiley-Interscience [John Wiley & Sons], New York (1978)
- [7] Ionescu, P.: *On manifolds of small degree*. *arXiv:math/0306205* (2003)
- [8] Jouanolou, J.P.: *Théorèmes de Bertini et applications*. *Progress in Mathematics*, 42. Birkhäuser Boston, Inc., Boston, MA (1983)
- [9] Mumford, D.: *Pathologies*. III. *Amer. J. Math.* **89**, 94–104 (1967)

JOSÉ CARLOS SIERRA, DEPARTAMENTO DE ÁLGEBRA, FACULTAD DE CIENCIAS MATEMÁTICAS, UNIVERSIDAD COMPLUTENSE DE MADRID, 28040 MADRID, SPAIN. E-MAIL jcsierra@mat.ucm.es